

DONNA

# Abdomen, Flanks (Love Handles), and Bra Fat



Before



After

19 weeks after two CoolSculpting® Elite treatments on her abdomen, and 14 weeks after three treatments on her flanks and bra fat. 8 weeks after eight CoolTone® treatments on her abdomen. **Results may vary.** The safety and efficacy of CoolSculpting® and CoolTone® in combination have not been evaluated by the FDA.

### CoolSculpting® and CoolSculpting® Elite Uses and Important Safety Information Uses

CoolSculpting® and CoolSculpting® Elite are FDA-cleared for the treatment of visible fat bulges in the submental (under the chin) and submandibular (under the jawline) areas, thigh, abdomen, and flank, along with bra fat, back fat, underneath the buttocks (also known as banana roll), and upper arm. It is also FDA-cleared to affect the appearance of lax tissue with submental area treatments. CoolSculpting® and CoolSculpting® Elite are not treatments for weight loss.

### Important Safety Information

This procedure is not for everyone. You should not be treated with CoolSculpting® or CoolSculpting® Elite if you suffer from cryoglobulinemia, cold agglutinin disease, or paroxysmal cold hemoglobinuria.

Tell your doctor if you are pregnant or have any medical conditions including recent surgery, pre-existing hernia, and any known sensitivities or allergies.

During the procedure you may experience sensations of pulling, tugging, mild pinching, intense cold, tingling, stinging, aching, cramping and discomfort at the treatment site. These sensations lessen as the area becomes numb. Following the procedure, typical side effects include temporary redness, swelling, blanching, bruising, firmness, stinging, tenderness, cramping, skin inflammation and aching. Sensory alteration (itching, skin sensitivity, tingling, and numbness) can persist up to several weeks after treatment. You may also experience throat discomfort/soreness and sensation of fullness in the back of the throat after submental or submandibular area treatment.

Rare side effects may happen in 1 to 10 out of 10,000 CoolSculpting® and CoolSculpting® Elite treatments (between 0.01% to 0.1%). One such rare side effect is a visible enlargement in the treated area, which may develop 2 to 5 months after treatment, will not resolve on its own, and may require surgical intervention for correction.

Please see full Important Safety Information for CoolSculpting® and CoolSculpting® Elite on CoolSculpting.com.

### Uses and Important Safety Information for CoolTone® Uses

The CoolTone® device is FDA-cleared for improvement of abdominal tone, strengthening of the abdominal muscles, and development for firmer abdomen. CoolTone® is also FDA-cleared for strengthening, toning, and firming of buttocks and thighs.

### Important Safety Information

The CoolTone® procedure is not for everyone. You should not have the CoolTone® treatment in areas with metal, active, electrical, or electronic implants/devices like cardiac pacemakers, implanted hearing devices, implanted defibrillators, implanted neurostimulators, drug pumps, or hearing aids.

Tell your doctor if you have any medical conditions as CoolTone® should not be used over a menstruating uterus, over areas of the skin that lack normal sensation, in patients with fever, malignant tumor, hemorrhagic conditions, epilepsy, recent surgical procedure, pulmonary insufficiency, or pregnancy.

CoolTone® should be used with caution in patients with Graves' disease (an autoimmune disorder that causes overactive thyroid), women with an intrauterine device (IUD), hernia in or near treatment site, active bleeding disorders, or seizure disorders. Women with an IUD should monitor for symptoms such as severe abdominal pain, cramping, spotting, or heavy bleeding develop during or post-treatment and seek medical advice if these occur.

Women who are close to menstruation may find that it comes sooner, or cramping is increased or intensified with CoolTone® treatments, therefore, it is recommended to not undergo treatment during this time of the month.

CoolTone® should not be used in the heart or head areas, areas of new bone growth, over the carotid sinus nerves, or over the neck or mouth. CoolTone® should not be applied over swollen, infected, inflamed areas or skin eruptions. Caution should be used for patients with suspected or diagnosed heart problems.

Common side effects may include, but may not be limited to, muscular pain, temporary muscle spasm, temporary joint or tendon pain, and redness at or near the treatment site.

Ask your Healthcare Provider if CoolTone® is right for you.

Please see full Important Safety Information for additional information at [cooltone.com](https://cooltone.com).