



KENID

Abdomen, Flanks (Love Handles), Back Fat, and Bra Fat



Before



After

12 weeks after three CoolSculpting® Elite treatments on her abdomen, flanks, back fat, and bra fat. Kenid lost 12.4 lb due to diet and exercise. CoolSculpting® is not a weight loss procedure. **Results may vary.**



Before



After

12 weeks after three CoolSculpting® Elite treatments on her abdomen, flanks, back fat, and bra fat. Kenid lost 12.4 lb due to diet and exercise. CoolSculpting® is not a weight loss procedure. **Results may vary.**

CoolSculpting® and CoolSculpting® Elite Uses and Important Safety Information

CoolSculpting® and **CoolSculpting® Elite** are FDA-cleared for the treatment of visible fat bulges in the submental (under the chin) and submandibular (under the jawline) areas, thigh, abdomen, and flank, along with bra fat, back fat, underneath the buttocks (also known as banana roll), and upper arm. It is also FDA-cleared to affect the appearance of lax tissue with submental area treatments. **CoolSculpting®** and **CoolSculpting® Elite** are not treatments for weight loss.

Important Safety Information

This procedure is not for everyone. You should not be treated with **CoolSculpting®** or **CoolSculpting® Elite** if you suffer from cryoglobulinemia, cold agglutinin disease, or paroxysmal cold hemoglobinuria.

Tell your doctor if you are pregnant or have any medical conditions including recent surgery, pre-existing hernia, and any known sensitivities or allergies.

During the procedure you may experience sensations of pulling, tugging, mild pinching, intense cold, tingling, stinging, aching, cramping and discomfort at the treatment site. These sensations lessen as the area becomes numb. Following the procedure, typical side effects include temporary redness, swelling, blanching, bruising, firmness, stinging, tenderness, cramping, skin inflammation and aching. Sensory alteration (itching, skin sensitivity, tingling, and numbness) can persist up to several weeks after treatment. You may also experience throat discomfort/soreness and sensation of fullness in the back of the throat after submental or submandibular area treatment.

Rare side effects may happen in 1 to 10 out of 10,000 **CoolSculpting®** and **CoolSculpting® Elite** treatments (between 0.01% to 0.1%). One such rare side effect is a visible enlargement in the treated area, which may develop 2 to 5 months after treatment, will not resolve on its own, and may require surgical intervention for correction.

Please see full Important Safety Information for **CoolSculpting®** and **CoolSculpting® Elite** on CoolSculpting.com.